

The 62nd William Lowell Putnam Mathematical Competition
Saturday, December 1, 2001

A-1 Consider a set S and a binary operation $*$, i.e., for each $a, b \in S$, $a * b \in S$. Assume $(a * b) * a = b$ for all $a, b \in S$. Prove that $a * (b * a) = b$ for all $a, b \in S$.

A-2 You have coins C_1, C_2, \dots, C_n . For each k , C_k is biased so that, when tossed, it has probability $1/(2k+1)$ of falling heads. If the n coins are tossed, what is the probability that the number of heads is odd? Express the answer as a rational function of n .

A-3 For each integer m , consider the polynomial

$$P_m(x) = x^4 - (2m+4)x^2 + (m-2)^2.$$

For what values of m is $P_m(x)$ the product of two non-constant polynomials with integer coefficients?

A-4 Triangle ABC has an area 1. Points E, F, G lie, respectively, on sides BC, CA, AB such that AE bisects BF at point R , BF bisects CG at point S , and CG bisects AE at point T . Find the area of the triangle RST .

A-5 Prove that there are unique positive integers a, n such that $a^{n+1} - (a+1)^n = 2001$.

A-6 Can an arc of a parabola inside a circle of radius 1 have a length greater than 4?

B-1 Let n be an even positive integer. Write the numbers $1, 2, \dots, n^2$ in the squares of an $n \times n$ grid so that the k -th row, from left to right, is

$$(k-1)n+1, (k-1)n+2, \dots, (k-1)n+n.$$

Color the squares of the grid so that half of the squares in each row and in each column are red and the other half are black (a checkerboard coloring is one possibility). Prove that for each coloring, the sum of the numbers on the red squares is equal to the sum of the numbers on the black squares.

B-2 Find all pairs of real numbers (x, y) satisfying the system of equations

$$\frac{1}{x} + \frac{1}{2y} = (x^2 + 3y^2)(3x^2 + y^2)$$

$$\frac{1}{x} - \frac{1}{2y} = 2(y^4 - x^4).$$

B-3 For any positive integer n , let $\langle n \rangle$ denote the closest integer to \sqrt{n} . Evaluate

$$\sum_{n=1}^{\infty} \frac{2^{\langle n \rangle} + 2^{-\langle n \rangle}}{2^n}.$$

B-4 Let S denote the set of rational numbers different from $\{-1, 0, 1\}$. Define $f : S \rightarrow S$ by $f(x) = x - 1/x$. Prove or disprove that

$$\bigcap_{n=1}^{\infty} f^{(n)}(S) = \emptyset,$$

where $f^{(n)}$ denotes f composed with itself n times.

B-5 Let a and b be real numbers in the interval $(0, 1/2)$, and let g be a continuous real-valued function such that $g(g(x)) = ag(x) + bx$ for all real x . Prove that $g(x) = cx$ for some constant c .

B-6 Assume that $(a_n)_{n \geq 1}$ is an increasing sequence of positive real numbers such that $\lim a_n/n = 0$. Must there exist infinitely many positive integers n such that $a_{n-i} + a_{n+i} < 2a_n$ for $i = 1, 2, \dots, n-1$?